## NOTE V.

## ON A NEW TAPHOZOUS FROM JAVA

BY

## Dr. F. A. JENTINK.

March 1907.

Dr. P. N. van Kampen offered some bats to our Museum, all from the neighborhood of Batavia, Java. Among them is a Taphozous, an adult male, with a radiometacarpal pouch and a gular sac. Among the very few Taphozous-species known from the East Indian Region, there only is a single one, presenting these two characteristics combined, viz. Taphozous longimanus Hardwicke, from the Indian Continent. Dobson (Catalogue, p. 385) described a variety from Flores under the name of Taphozous longimanus leucopleurus (by an error in Trouessart's splendid ,,Catalogus Mammalium" mentioned as leucopteurus and leucopterus). I registered in the Notes from the Leyden Museum, 1897, p. 54, two specimens, collected by Mr. Goedhuis at Sintang, Borneo, a female and a young male, under the specific title of Taphozous longimanus Hardw., the first known specimens of the Taphozous longimanus-group from Borneo, meanwhile Oldfield Thomas (Ann. Mag. Nat. Hist. 1898, II, p. 246) described another specimen, an adult female, collected by Mr. Everett in 1892 at Labuan, Borneo, as a subspecies, Taphozous longimanus albipinnis. Up to the present no specimens of the longimanus-group have been recorded from Java; therefore it prima facie seems very probable that our Batavia-specimen may turn out to be another form of the rare and highly interesting longimanus-group.

Notes from the Leyden Museum, Vol. XXIX.

I describe it, in honour of its Donor, under the name of Taphozous Kampenii, n. sp.

Some measurements (in mM.) of the type, an adult male in spirits, compared with longimanus (Dobs. Cat.) from the Indian Continent; leucopleurus (Dobs. Cat.) from Flores; the adult female from Sintang in the Leyden Museum; Oldfield Thomas' type from Labuan, T. l. albipinnis, besides a specimen in the British Museum also from Borneo, Sarawak. The measurements of the two latter specimens I thankfully owe to the kindness of Mr. Knud Andersen.

1	Batavia.	1nd. Cont.	Flores.	Sintang.	Labuan.	Sarawak.
tail	. 21	28.75	27.5	23	20.2	19.5
ear	. 18	20	20	18	18.8	18.4
forearm	. 57	61.25	57.5	57.5	56	
third finger, metacarp	. 58	)	57.5	55		
" " 1st phalanx.	. 23	105	22.5	20		
" " 2nd " .			25	21		
fifth " metacarp	. 34	1	33.75	33		
" " 1st phalanx.	. 12	53.75	12.5	11		
" " 2nd "			10	9		
tibia	. 24	23.75	22.5	23		
foot	. 12.5	11.25	12.5	11		
calcaneum	. 16	21.25		15		

As the material is very limited — 1 specimen from Batavia, 1 from Flores and 3 from Borneo — the above given measurements are by no means decisive; it at present cannot made out whether there is a single very variable species, represented in different localities by distinct forms, or if perhaps Borneo, Java, Flores a. s. o. are inhabited by sharply circumscribed distinct species? It is very likely that our Sintang-form is what Thomas described as albipinnis and that the Batavia-specimen is in reality different; the more as I observed in it some characteristics different from those in our Sintang-specimen, which I shortly will point out.

Dobson said "that the fur in the specimens of longima-Notes from the Leyden Museum, Vol. XXIX. nus varies very much", however it seems that the continental individuals are always darker (reddish-brown or black) colored than the Borneo-representatives; moreover the latter are white-winged; so our Sintang-female too, in striking contrast with our Java-male, with much darker fur and sooty colored wings. In the Java-individual the nasal apertures seem closer together than in the Sintang-specimen, meanwhile there is a much deeper groove between them in the Java-one. The ear of the Batavia-male, although of the same length as the ear of the Sintang-female, is somewhat broader, and the outer margin of the ear-conch presents a slight excavation in the former, being smooth in the Sintang-specimen. Very small seems the tail of the Borneo- and Java-specimen compared with the same organ of the Continental- as well as of the Flores-form. As I remarked, the material at hand is by no means sufficient, however the short hinds given above are very characteristic and will enable to recognize our Kampenii from Batavia.